

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/00		A2	(11) International Publication Number: WO 99/09967
			(43) International Publication Date: 4 March 1999 (04.03.99)
(21) International Application Number: PCT/US98/17459		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HR, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 21 August 1998 (21.08.98)			
(30) Priority Data: 60/056,966 26 August 1997 (26.08.97) US 60/076,859 5 March 1998 (05.03.98) US 60/085,360 13 May 1998 (13.05.98) US 9814415.7 3 July 1998 (03.07.98) GB 9815851.2 21 July 1998 (21.07.98) GB			
(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		Published Without international search report and to be republished upon receipt of that report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): WHITNEY, Edwin, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MANTELL, Geraldine [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			
(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			
(54) Title: CHOLESTEROL-LOWERING THERAPY			
(57) Abstract <p>Methods are described for preventing or reducing the risk of a first occurrence of a cardiovascular event using an HMG-CoA reductase inhibitor alone or in combination with another lipid altering agent. Subjects to be treated are those having an average serum total cholesterol level, an average to mildly elevated serum low-density lipoprotein cholesterol level, and a below average serum high-density lipoprotein cholesterol level, with no history of clinically evident coronary disease.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

CHOLESTEROL-LOWERING THERAPY

FIELD OF THE INVENTION

5 The instant invention involves a method of using a
cholesterol reducing agent such as a 3-hydroxy-3-methylglutaryl
coenzyme A (HMG-CoA) reductase inhibitor (or HMG-CoA RI) alone or
in combination with other lipid altering agents for preventing or
reducing the risk of first occurrence of a cardiovascular event.

10

BACKGROUND OF THE INVENTION

 It has been clear for several decades that elevated blood
cholesterol is a major risk factor for coronary heart disease (CHD), and
many studies have shown that the risk of CHD events can be reduced by
15 lipid-lowering therapy. Prior to 1987, the lipid-lowering
armamentarium was limited essentially to a low saturated fat and
cholesterol diet, the bile acid sequestrants (cholestyramine and
colestipol), nicotinic acid (niacin), the fibrates and probucol.
Unfortunately, all of these treatments have limited efficacy or
20 tolerability, or both. With the introduction of lovastatin, the first
inhibitor of HMG-CoA reductase to become available for prescription in
1987, for the first time physicians were able to obtain comparatively large
reductions in plasma cholesterol with very few adverse effects.

 Recent studies have unequivocally demonstrated that
25 lovastatin, simvastatin, and pravastatin, which are members of the
HMG-CoA reductase inhibitor class, slow the progression of
atherosclerotic lesions in the coronary and carotid arteries. Simvastatin
and pravastatin have also been shown to reduce the risk of coronary
heart disease events in patients with hypercholesterolemia and/or CHD,
30 and in the case of simvastatin a highly significant reduction in the risk
of coronary death and total mortality has been shown by the
Scandinavian Simvastatin Survival Study. This study also provided
evidence for a reduction in cerebrovascular events. Additional studies
have shown that HMG CoA RI's may have an effect on platelet
35 aggregation.

SUMMARY OF THE INVENTION

One object of the instant invention is to provide a novel method for preventing or reducing the risk of a first occurrence of a cardiovascular event in a subject having an average to mildly elevated level of LDL cholesterol, and below average high-density lipoprotein (HDL) cholesterol, with no clinical evidence of coronary heart disease, comprising administering a prophylactically effective amount of a lipid altering agent such as an HMG-CoA reductase inhibitor either alone or in combination with another lipid altering agent such as a fibrate, or niacin to the subject. Additional objects will be evident from the following detailed description.

DETAILED DESCRIPTION OF THE INVENTION

Examples of HMG-CoA reductase inhibitors that may be used include but are not limited to lovastatin (MEVACOR®; see US Patent No. 4,231,938), simvastatin (ZOCOR®; see US Patent No. 4,444,784), pravastatin (PRAVACHOL®; see US Patent No. 4,346,227), fluvastatin (LESCOL®; see US Patent No. 5,354,772), atorvastatin (LIPITOR®; see US Patent No. 5,273,995) and cerivastatin (also known as rivastatin; see US Patent No. 5,177,080), and the pharmaceutically acceptable salt, ester and lactone forms thereof. The structural formulas of these and additional HMG-CoA reductase inhibitors that may be used in the instant methods are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs", *Chemistry & Industry*, pp. 85-89 (5 February 1996). The HMG-CoA RI is selected from lovastatin, cerivastatin, atorvastatin, pravastatin and simvastatin, a sub-class comprises simvastatin and lovastatin. Examples of fibrates that may be used in combination with the HMG-CoA RI include, but are not limited to benzofibrate, ciprofibrate, fenofibrate, gemfibrozil and clofibrate.

The term HMG-CoA reductase inhibitor is intended to include all pharmaceutically acceptable salt, ester and lactone forms of compounds which have HMG-CoA reductase inhibitory activity, and therefore the use of such salts, esters and lactone forms is included within the scope of this invention.

Compounds which have inhibitory activity for HMG-CoA reductase can be readily identified by using assays well-known in the art. For example, see the assays described or cited in U.S. Patent 4,231,938 at col. 6, and WO 84/02131 at pp. 30-33.

5 Ester derivatives of the described compounds may act as prodrugs which, when absorbed into the bloodstream of a warm-blooded animal, may cleave in such a manner as to release the drug form and permit the drug to afford improved therapeutic efficacy.

 Herein, the term "pharmaceutically acceptable salts" shall
10 mean non-toxic salts of the compounds employed in this invention which are generally prepared by reacting the free acid with a suitable organic or inorganic base. Examples of such salts include, but are not limited to, acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium, calcium edetate, camsylate, carbonate,
15 chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycollylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynapthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate,
20 methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, pamaote, palmitate, panthothenate, phosphate/diphosphate, polygalacturonate, potassium, salicylate, sodium, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide, valerate.

25 Prophylactically effective amounts of the HMG-CoA reductase inhibitors are suitable for use in the compositions and methods of the present invention. The term "prophylactically effective amount" is intended to mean that amount of a pharmaceutical drug that will prevent or reduce the risk of occurrence of the biological or medical
30 event that is sought to be prevented of a tissue, a system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician. The dosage regimen utilizing an HMG-CoA RI is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the subject; the severity of the
35 condition to be treated; the route of administration; the renal and hepatic

function of the subject; and the particular compound or salt or ester thereof employed. A consideration of these factors is well within the purview of the ordinarily skilled clinician for the purpose of determining the prophylactically effective amounts of the drug needed to prevent or
5 reduce the risk of occurrence of the condition sought to be prevented. In one embodiment, the prophylactically effective amount to be used in the methods of this invention is that amount of drug sufficient to lower the human subject's LDL cholesterol to a target level of 130 mg/dl or less; in a sub-embodiment to a target level of 115 mg/dl or less; and in a further
10 sub-embodiment to a target level of 110 mg/dl or less. The ordinarily skilled clinician will be able to titrate the subject to the appropriate amount of lipid altering agent such as an HMG-CoA RI which, when taken on a daily basis, will allow the patient to reach this goal.

The term "patient" or "subject" includes mammals,
15 especially humans, who take a lipid altering agent for any of the uses described herein. Administration of the lipid altering agent to the subject includes both self-administration and administration to the subject by another person.

Dosage information for HMG-CoA RI's is well known in the
20 art, since several are marketed in the U.S. In particular, the daily dosage amounts of the HMG-CoA reductase inhibitor may be the same or similar to those amounts which are employed for anti-hypercholesterolemic treatment and which are described in the *Physicians' Desk Reference* (PDR). For example, see the 50th Ed. of the
25 PDR, 1996 (Medical Economics Co); in particular, see at page 216 the heading "Hypolipidemics," sub-heading "HMG-CoA Reductase Inhibitors," and the reference pages cited therein. Preferably, the oral dosage amount of HMG-CoA RI is from about 1 to 200 mg/day, and more preferably from about 5 to 160 mg/day. However, dosage amounts will
30 vary depending on the potency of the specific HMG-CoA reductase inhibitor used as well as other factors as noted above. An HMG-CoA RI which has sufficiently greater potency may be given in sub-milligram daily dosages.

As examples, the daily dosage amount for simvastatin may
35 be selected from 5 mg, 10 mg, 20 mg, 40 mg, and 80 mg; for lovastatin, 10

mg, 20 mg, 40 mg and 80 mg; for fluvastatin sodium, 20 mg, 40 mg and 80 mg; and for pravastatin sodium, 10 mg, 20 mg, and 40 mg. The daily dosage amount for cerivastatin may be in the range of from 0.1 mg to 0.8 mg, and more particularly from 0.2 mg to 0.8 mg, including dosage
5 amounts of 0.2 mg, 0.3 mg, 0.4 mg and 0.8 mg. Oral administration may be in single or divided doses of two, three, or four times daily, although a single daily dose of the HMG-CoA RI is preferred.

Pharmaceutical formulations for HMG-CoA reductase inhibitors are well-known to those skilled in the art, as evidenced by the
10 information provided in the 1996 PDR. While the HMG-CoA reductase inhibitor can be administered orally or parenterally, oral dosing is preferred.

For example, the active agents employed in the instant therapy can be administered in such oral forms as tablets, capsules
15 (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. Oral formulations are preferred. The instant invention includes the use of oral rapid-release as well as time-controlled release pharmaceutical formulations, particularly as
20 described in U.S. Patent No. 5,366,738.

The active drug can be administered in admixture with pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs,
25 syrups and the like, and consistent with conventional pharmaceutical practices. See, for example, *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with a non-toxic,
30 pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, modified sugars, modified starches, methyl cellulose and its derivatives, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and other reducing and non-reducing sugars, magnesium stearate, steric acid, sodium stearyl fumarate, glyceryl behenate,
35 calcium stearate and the like. For oral administration in liquid form,

the drug components can be combined with non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring and flavoring agents can also be
5 incorporated into the mixture. Stabilizing agents such as antioxidants (BHA, BHT, propyl gallate, sodium ascorbate, citric acid) can also be added to stabilize the dosage forms. Other suitable components include gelatin, sweeteners, natural and synthetic gums such as acacia, tragacanth or alginates, carboxymethylcellulose, polyethylene glycol,
10 waxes and the like.

The active drug can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol,
15 stearylamine or phosphatidylcholines.

Active drug may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. Active drug may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinyl-
20 pyrrolidone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxy-ethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, active drug may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid,
25 polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels.

The methods of the invention may be used to prevent or
30 reduce the risk of a first occurrence of a fatal or non-fatal cardiovascular event. The term "cardiovascular event" includes but is not limited to fatal and non-fatal acute major coronary events, coronary revascularization procedures, peripheral vascular disease, stable angina, and cerebrovascular insufficiency such as stroke.

The term "acute major coronary event" is intended to include fatal myocardial infarction; witnessed and unwitnessed sudden cardiac death and sudden death occurring from 1 hour up to 24 hours after collapse; non-fatal myocardial infarction including definite acute Q-wave myocardial infarction, non-Q-wave myocardial infarction, and silent subclinical (remote) myocardial infarction; and unstable angina pectoris. As used herein, the term "myocardial infarction" is intended to include both Q-wave and non-Q-wave myocardial infarction and silent subclinical (remote) myocardial infarction.

Subjects to be treated with the instant methods are those having an average to mildly elevated serum total cholesterol level, which is intended herein to be a level less than or equal to about 260 mg/dl. In particular, the patients to be treated may have a serum total cholesterol level of less than or equal to 240 mg/dl, and more particularly under 200 mg/dl. For example, the subjects to be treated may have a serum total cholesterol level from 180 mg/dl to 264 mg/dl. In one embodiment, the subjects to be treated have a below average high-density lipoprotein cholesterol level, i.e., less than or equal to 50 mg/dl, and an average to mildly elevated low-density lipoprotein cholesterol level, i.e., less than or equal to 190 mg/dl, prior to treatment. In another embodiment the low-density lipoprotein cholesterol level is 130 mg/dl to 190 mg/dl prior to treatment. In a still further embodiment, the LDL-cholesterol is < 130 mg/dl if the ratio of total cholesterol to HDL-cholesterol is > 6. The subjects to be treated also have no clinically evident coronary heart disease.

In accordance with this invention, a prophylactically effective amount of an HMG-CoA RI or a combination with another lipid altering agent can be used for the preparation of a medicament useful for preventing or reducing the risk of a first occurrence of a cardiovascular event in a subject having an average to mildly elevated level of serum total cholesterol, an average to mildly elevated level of low-density lipoprotein cholesterol, and a below average level of high-density lipoprotein cholesterol.

The following is a description of a clinical trial referred to herein as "AFCAPS/TexCAPS" or "AFCAPS" employing lovastatin to exemplify the methods of the present invention.

5

EXAMPLE 1

AFCAPS/TexCAPS was a primary prevention endpoint event trial which: (1) included unstable angina as a primary endpoint, reflecting the trend to treat coronary heart disease aggressively before a myocardial infarction has occurred; 2) involved aggressive pharmacologic intervention, with titration, to target an LDL-cholesterol goal lower than current National Cholesterol Education Panel guidelines for primary prevention; and 3) included a cohort comprised of men and women with a broad age range encompassing the elderly and included persons with a lipid profile consisting of average total and LDL-cholesterol, below average HDL-cholesterol levels; total cholesterol/HDL-cholesterol ratio greater than or equal to 4.5, with a baseline mean ratio of 6.1 for the cohort; and LDL/HDL-cholesterol ratio greater than or equal to 4.

Entrance inclusion criteria were LDL-cholesterol 130-190 mg/dl (or < 130 if the ratio of total cholesterol/HDL is > 6) and HDL-cholesterol \leq 45 mg/dl for men and \leq 47 mg/dl for women.

The trial was designed to study the effect of LDL-cholesterol reduction in a cohort with average to mildly elevated LDL-cholesterol and a below average HDL-cholesterol. National Cholesterol Education Panel guidelines for adults (see National Cholesterol Education Program Expert Panel, Summary of the second report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II), *JAMA*, 1993;269:3015-3023) already incorporate HDL-cholesterol concentration in recommendations for estimating CHD risk and selecting lipid-altering therapy.

35

STUDY DESIGN

AFCAPS/TexCAPS was a double-blind, randomized, placebo controlled trial designed and powered to investigate whether chronic lipid-lowering, with lovastatin will decrease the rate of first acute major coronary events (e.g. sudden cardiac death, fatal and non-fatal myocardial infarction and unstable angina) compared to intervention with diet alone during at least 4.5 years of follow-up in a cohort without clinical evidence of atherosclerotic cardiovascular disease and with average to mildly elevated total cholesterol, average to mildly elevated LDL-cholesterol, and below average HDL-cholesterol. Secondary objectives included whether chronic treatment with lovastatin, compared to placebo, would decrease cardiovascular morbidity and mortality across the spectrum of clinical events, by measuring the rates of: (1) fatal and non-fatal coronary revascularization procedures (2) unstable angina, (3) fatal and non-fatal myocardial infarction, (4) fatal and non-fatal cardiovascular events, (5) fatal and non-fatal coronary events.

All participants underwent dietitian taught group instruction in the American Heart Association Step 1 diet commencing at least 12 weeks prior to randomization (Week -12) and reinforced at Weeks -4 and -2. Baseline observations included collection of medical history, health habit and demographic data, physical exam, eye exam (San Antonio clinic only), chest X-ray, mammogram (women), electrocardiogram and analysis of lipids/lipoproteins, hematology, blood chemistry and urine. A 2-week placebo baseline run-in was followed by randomization of eligible compliant participants to treatment with lovastatin or placebo.

Participants included 6605 men and women with normal to mildly elevated total and LDL-cholesterol and below average HDL-cholesterol defined as: total cholesterol 180-264 mg/dl, LDL-cholesterol 130-190 mg/dl, HDL-cholesterol ≤ 45 mg/dl for men and ≤ 47 mg/dl for women and triglycerides ≤ 400 mg/dl. All lipid/lipoprotein entrance criteria had to be met on average at both four and two weeks prior to randomization (after at least eight weeks of diet therapy). In addition,

for both total and LDL-cholesterol, the two eligibility determinations had to be within 15% of each other.

Excluded for clinical evidence of atherosclerotic cardiovascular disease were men and women who had: prior history of myocardial infarction; angina; claudication; cerebrovascular accident; or transient ischemic attack. Also excluded were those aged > 73 years or < 45 years (men) or < 55 years (women) or those with secondary hyperlipoproteinemia, nephrotic syndrome, uncontrolled or insulin-dependent diabetes mellitus or uncontrolled hypertension. At their discretion, investigators also could have excluded those who would have difficulty completing a study of at least 5 years duration (e.g. due to compliance problems or reduced life expectancy).

Baseline risk factors in the tested population were as follows:

Family history of premature coronary artery disease: 16%;
Active cigarette smoker: 12%;
Hypertension: 22%;
Diabetes: 2%.
HDL < 35 mg/dL: 35%

Participants were randomized to either placebo or lovastatin 20 mg once a day and were stratified by center and gender so that there was an equal chance of being assigned to lovastatin or placebo. Formal procedures ensured maintenance of the study blind. Participants, investigators, Steering Committee members and those providing participant care, monitoring or managing data, or adjudicating endpoints were blinded. For participant safety, an independent Data and Safety Monitoring Board was not blinded. In addition, two programmers and the study statistician who performed analyses for the safety monitoring board had access to either the allocation schedule or to the unblinded lipid value.

Baseline measurements on the day of randomization included lipid analysis (including Apo A1 and Apo B), hematology, blood chemistry and urinalysis.

During the first 48 weeks of active treatment, participants returned to clinic at 6 week intervals. At each visit, participants were

asked about adverse events and underwent laboratory safety tests for liver enzymes and creatine kinase. Every 12 weeks, participants also receive dietary reinforcement. During the first 6 months, a lipid profile was performed at every visit. At Week 48, participants received dietary advice and underwent an extensive evaluation that included the tests routinely done at each visit, as well as physical exam, electrocardiogram, mammography (women), ophthalmological examination (San Antonio clinic only), complete blood chemistry, hematology and urinalysis.

After Week 48, the interval between visits was longer (Week 60, Month 18, then every 6 months). At week 60 and at all "mid-year" visits, adverse experience inquiry and laboratory safety tests were performed. The year-end visit included all tests and observations described at the Week 48 visit. At Week 48 and Year 5, Apo A1 and Apo B were measured. At all protocol visits, pharmacists assessed compliance by pill counts.

To attempt to achieve an on-treatment LDL-cholesterol of 110 mg/dl or lower, participants in the lovastatin treatment group could be titrated to lovastatin 40 mg once a day based upon LDL-cholesterol values at 6 and 12 weeks after randomization. Participants in the lovastatin treatment group with LDL-cholesterol > 110 mg/dl at both visits were titrated at week 18 to 40 mg once a day of lovastatin by taking two 20 mg tablets once a day. To maintain the blind, an equal number of randomly selected participants in the placebo treatment group were also titrated to two tablets per day. To maintain the blind, an equal number from the titrated placebo group also had their doses reduced.

Participants were flagged for withdrawal by the unblinded programmer, if LDL-cholesterol values > 195 mg/dl on successive visits following titration. The investigator, the participant and those involved in the participant's care were not unblinded to treatment group or during double blind.

The primary endpoint events included sudden cardiac death, fatal and non-fatal myocardial infarction and unstable angina. They are defined as follows:

I. Fatal Myocardial Infarction or Sudden Cardiac Death

The definition requires that there be no non-cardiac cause of death and one of the following:

- Fatal Myocardial Infarction - death within 28 days from the onset of symptoms of a definite acute myocardial infarction
- Witnessed Unexpected Sudden Cardiac Death - within 1 hour of symptoms
- Death occurring >1 hour but <24 hours after collapse
- Unwitnessed Unexpected Death, Presumed Sudden - must have confirming autopsy data or, if autopsy not performed, preceding history of CHD events or symptoms

II. Non-Fatal Myocardial Infarction

- Acute Q-Wave Myocardial Infarction - requires definitive ECG
- Acute Non-Q-Wave Myocardial Infarction - definitive ECG or, if equivocal, enzymes must be diagnostic. Non-Q-Wave Myocardial Infarction includes myocardial infarctions reperfused by either mechanical or pharmacologic means providing there is supporting ECGs and enzyme data
- Silent Subclinical (Remote) Myocardial Infarction - definitive ECG, or, if ECG is equivocal, focal wall motion abnormality consistent with myocardial infarction on rest echo or stress thallium (fixed defect) and on catheterization, a $\geq 50\%$ stenosis in a major corresponding epicardial vessel. Participants who have had a cardiac catheterization as the first diagnostic test for presumed silent (or remote) myocardial infarction are considered to have met criteria for an endpoint event if the catheterization findings indicate focal wall abnormalities consistent with myocardial infarction and 50% stenosis in a corresponding artery

III. Unstable Angina

New onset exertional and/or accelerated or rest angina and requires at least one of the following:

- Stress perfusion study: 1mm ST segment changes and reversible defect
- 90% epicardial vessel stenosis or 50% stenosis in the Left Main

- ≥ 1 mm ST segment changes with pain on stress testing and/or resting ECG and evidence of 50% stenosis in a major epicardial vessel

5 Note that 'angina' was adjudicated as a secondary endpoint event if, prior to the hospitalization, the participant was asymptomatic for > 2 weeks or has been stable for > 1 month (defined as 28 days) even if the criteria for 'unstable angina', noted above, were met.

All subjects were followed until the decision to end the study after a median duration of 5.2 years of treatment.

10 The trial design for the final analysis provided 90 to 97% power, respectively, to detect the reductions in the number of patients experiencing as a first event any of those shown in the following Table 1:

TABLE 1 - EFFICACY ENDPOINTS

	Risk Reduction	Relative Risk (C.I.)
ENDPOINTS		
Primary Endpoint: acute major coronary events defined as fatal and non-fatal myocardial infarction, unstable angina or sudden cardiac death	37%	0.63 (0.50, 0.79), p=0.00008
Secondary Endpoints		
• Revascularizations	33%	0.67 (0.52, 0.85), p=0.001
• Unstable Angina	32%	0.68 (0.48, 0.95), p=0.023
• Fatal and Nonfatal MI	40%	0.60 (0.43, 0.83), p=0.002
• Fatal and Nonfatal Cardiovascular Events	25%	0.75 (0.62, 0.91), p=0.003
• Fatal and Nonfatal Coronary Events	25%	0.75 (0.61, 0.92), p=0.006

Thus, the risk of an acute major coronary event using
5 lovastatin was reduced by at least about 20% and more particularly
about 21% to 50%, and illustratively about 30% to 40%. The risk
reduction of an acute major coronary event with simvastatin,
pravastatin, fluvastatin, cerivastatin, or atorvastatin is also expected
to be at least about 20%, and more particularly about 21% to 50%, and
10 illustratively about 30% to 40%.

Similarly, the risk of a fatal and non-fatal
cardiovascular event using lovastatin was reduced by at least about
9% and more particularly about 9% to 38%, and illustratively about

20% to 30%. The risk reduction of a fatal or non-fatal cardiovascular event with simvastatin, pravastatin, fluvastatin, cerivastatin or atorvastatin is also expected to be at least about 9%, and more particularly about 9% to 38%, and illustratively about 20% to 30%.

5 Also, the risk of unstable angina using lovastatin was reduced by at least about 5% and more particularly about 5% to 52%, and illustratively about 25% to 40%. The risk reduction of unstable angina with simvastatin, pravastatin, fluvastatin, cerivastatin, or atorvastatin is also expected to be at least about 5%, and more
10 particularly about 5% to 52%, and illustratively about 25% to 40%.

Also, the risk of revascularization using lovastatin was reduced by at least about 15% and more particularly about 15% to 48%, and illustratively about 30% to 40%. The risk reduction of revascularization with simvastatin, pravastatin, fluvastatin,
15 cerivastatin, or atorvastatin is also expected to be at least about 15%, and more particularly about 15% to 48%, and illustratively about 30% to 40%.

Similarly, the risk of a fatal or non-fatal myocardial infarction using lovastatin was reduced by at least about 17% and
20 more particularly about 17-57%, and illustratively, about 30-50%. The risk reduction of a fatal or non-fatal myocardial infarction using simvastatin, pravastatin, fluvastatin, cerivastatin or atorvastatin is also expected to be at least about 17%, and more particularly about 17-57%, and illustratively, about 30-50%.

25 To further evaluate the effect of treatment, risk reduction for first acute major coronary events (the primary endpoint) was investigated by LDL-cholesterol (LDL-C) and HDL-cholesterol (HDL-C) tertiles at the time of randomization. The results are shown in the following Table 2 (with 95% confidence interval, or 95 % C.I.), where “%
30 change” indicates the change in LDL-C or HDL-C, as appropriate, from baseline to the one year point:

TABLE 2

	% change	Risk Reduction	Relative Risk (95% C.I.)
LDL-C (mg/dl)			
< 142	-20.1	34%	0.66 (0.43, 1.00)
142.0 - 156.8	-26.0	36%	0.64 (0.42, 0.99)
> 156.9	-28.9	41%	0.59 (0.41, 0.85)
HDL-C (mg/dl)			
< 34.0	7.4	45%	0.55 (0.38, 0.82)
34.5 - 39.5	6.2	43%	0.57 (0.39, 0.84)
> 39.5	4.3	15%	0.85 (0.54, 1.32)

5 The risk reduction achieved with lovastatin was independent of baseline LDL-C and HDL-C, thus demonstrating that benefit is conferred even in those subjects in the lowest LDL-C tertile. Similar results are also expected to be achieved with simvastatin, pravastatin, fluvastatin, cerivastatin and atorvastatin.

10 While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the invention. For example, effective dosages other than the particular dosages as set forth herein above may be
15 applicable as a consequence of variations in the responsiveness of the mammal being treated for any of the indications for the active agents used in the instant invention as indicated above. Likewise, the specific pharmacological responses observed may vary according to and depending upon the particular active compound selected or
20 whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It

is intended, therefore, that the invention be defined by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.

WHAT IS CLAIMED IS:

1. A method for preventing or reducing the risk of a first occurrence of a cardiovascular event comprising administering a prophylactically effective amount of an HMG-CoA reductase inhibitor alone or in combination with another lipid altering agent to a subject having a serum high-density lipoprotein cholesterol level less than or equal to 50 mg/dl prior to treatment.
2. The method of Claim 1 wherein the subject has an average level of serum low-density lipoprotein cholesterol prior to treatment.
3. The method of Claim 2 wherein the subject has a low-density lipoprotein cholesterol level less than or equal to 190 mg/dl prior to treatment.
4. The method of Claim 1 wherein the subject has an average level of serum total cholesterol prior to treatment.
5. The method of claim 1 wherein the subject has a serum triglycerides level of less than or equal to 400 mg/dl prior to treatment.
6. The method of claim 1 wherein the subject has a ratio of total cholesterol/HDL-cholesterol of greater than or equal to 4.5.
7. The method of claim 1 wherein the subject has no history of clinically evident coronary heart disease.
8. The method of Claim 7 wherein the subject has an average level of serum low-density lipoprotein cholesterol and an average level of serum total cholesterol prior to treatment.

9. The method of claim 8 wherein the subject has a ratio of total cholesterol/HDL-cholesterol of greater than or equal to 4.5.

10. The method of claim 9 wherein the subject has a serum triglycerides level of less than or equal to 400 mg/dl prior to treatment.

11. The method of Claim 1 wherein the prophylactically effective amount of the HMG-CoA reductase inhibitor is an amount sufficient to lower the subject's low-density lipoprotein cholesterol level to 130 mg/dl or less.

12. The method of Claim 11 wherein the prophylactically effective amount of the HMG-CoA reductase inhibitor is an amount sufficient to lower the treated subject's low-density lipoprotein cholesterol level to 115 mg/dl or less.

13. The method of Claim 12 wherein the prophylactically effective amount of the HMG-CoA reductase inhibitor is an amount sufficient to lower the treated subject's low-density lipoprotein cholesterol level to 110 mg/dl or less.

14. The method of Claim 1 wherein the subject has a serum total cholesterol level less than or equal to 260 mg/dl.

15. The method of Claim 1 wherein the subject is female and at least 55 years old and has a serum high-density lipoprotein cholesterol level less than or equal to 47 mg/dl prior to treatment.

16. The method of Claim 1 wherein the subject is male and at least 45 years old and has a serum high-density lipoprotein cholesterol level less than or equal to 45 mg/dl prior to treatment.

17. The method of Claim 1 wherein the reduction of risk of a first occurrence of a cardiovascular event is at least about 10%.

18. The method of Claim 17 wherein the reduction of risk of a first occurrence of a cardiovascular event is at least about 20-30%

5 19. The method of Claim 1 wherein the subject is administered a prophylactically effective amount of an HMG-CoA reductase inhibitor regularly for at least 1 year.

20. The method of Claim 19 wherein the subject is
10 administered a prophylactically effective amount of an HMG-CoA reductase inhibitor regularly for at least 2 years.

21. The method of Claim 20 wherein the subject is
15 administered a prophylactically effective amount of an HMG-CoA reductase inhibitor regularly for at least 4.5 years.

22. The method of Claim 1 wherein the cardiovascular event is selected from the group consisting of fatal and non-fatal acute major coronary events, coronary revascularization procedures,
20 peripheral vascular disease, stable angina, and cerebrovascular accidents.

23. The method of Claim 22 wherein the cardiovascular event is selected from the group consisting of fatal and non-fatal acute
25 major coronary events and coronary revascularization procedures.

24. The method of Claim 23 wherein the fatal and non-fatal acute major coronary events are selected from the group consisting of fatal myocardial infarction; witnessed and unwitnessed sudden
30 cardiac death and sudden death occurring from 1 hour up to 24 hours after collapse; non-fatal myocardial infarction; definite acute Q-wave myocardial infarction; non-Q-wave myocardial infarction; silent subclinical (remote) myocardial infarction; and unstable angina pectoris.

35

25. The method of Claim 1 wherein the HMG-CoA reductase inhibitor is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin, cerivastatin and the pharmaceutically acceptable salt, ester and lactone forms thereof.

5

26. The method of Claim 25 wherein the HMG-CoA reductase inhibitor is selected from the group consisting of lovastatin and simvastatin.

10

27. A method for preventing or reducing the risk of a first occurrence of a cardiovascular event by at least 20-30% comprising administering a prophylactically effective amount of an HMG-CoA reductase inhibitor to a subject having, prior to treatment, a serum high-density lipoprotein cholesterol level less than or equal to 50 mg/dl, a
15 serum low-density lipoprotein cholesterol level less than or equal to 190 mg/dl, a serum total cholesterol level less than or equal to 260 mg/dl, a ratio of total cholesterol/HDL-cholesterol of greater than or equal to 4.5, and no history of clinically evident coronary heart disease.

20

28. The method of Claim 27 wherein the prophylactically effective amount of the HMG-CoA reductase inhibitor is an amount sufficient to lower the subject's low-density lipoprotein cholesterol level to 130 mg/dl or less.

29. The method of Claim 28 wherein the prophylactically effective amount of the HMG-CoA reductase inhibitor is an amount sufficient to lower the subject's low-density lipoprotein cholesterol level to 115 mg/dl or less.

5

30. The method of Claim 29 wherein the prophylactically effective amount of the HMG-CoA reductase inhibitor is an amount sufficient to lower the subject's low-density lipoprotein cholesterol level to 110 mg/dl or less.

10

31. The method of Claim 27 wherein the cardiovascular event is selected from the group consisting of fatal and non-fatal acute major coronary events, coronary revascularization procedures, peripheral vascular disease, stable angina, and cerebrovascular accidents.

15

32. The method of Claim 31 wherein the cardiovascular event is selected from the group consisting of fatal and non-fatal acute major coronary events and coronary revascularization procedures.

20

33. A method of reducing the risk of an acute major coronary event or a coronary revascularization procedure in a human subject with average to mildly elevated LDL cholesterol, below average HDL cholesterol and no symptomatic cardiovascular disease which comprises the administration of a prophylactically effective amount of an HMG-CoA reductase inhibitor.

25

34. The method of Claim 33 wherein the acute coronary event is selected from myocardial infarction, unstable angina and sudden cardiac death.

30

35. The method of Claim 33 where the acute coronary event is selected from myocardial infarction and unstable angina.

36. The method of Claim 33 wherein the HMG-CoA reductase inhibitor is selected from lovastatin, simvastatin, fluvastatin, cerivastatin and atorvastatin.

5 37. The method of Claim 36 wherein the HMG-CoA reductase inhibitor is lovastatin or simvastatin.

38. The method of Claim 34 wherein the HMG-CoA reductase inhibitor is selected from lovastatin, simvastatin, fluvastatin,
10 cerivastatin and atorvastatin.

39. The method of Claim 38 wherein the HMG-CoA reductase inhibitor is lovastatin or simvastatin.

15

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/365, 31/40, 31/44		A3	(11) International Publication Number: WO 99/09967
			(43) International Publication Date: 4 March 1999 (04.03.99)
(21) International Application Number: PCT/US98/17459		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HR, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 21 August 1998 (21.08.98)			
(30) Priority Data: 60/056,966 26 August 1997 (26.08.97) US 60/076,859 5 March 1998 (05.03.98) US 60/085,360 13 May 1998 (13.05.98) US 9814415.7 3 July 1998 (03.07.98) GB 9815851.2 21 July 1998 (21.07.98) GB			
(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(72) Inventors; and (75) Inventors/Applicants (for US only): WHITNEY, Edwin, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MANTELL, Geraldine [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		(88) Date of publication of the international search report: 20 May 1999 (20.05.99)	
(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			
(54) Title: CHOLESTEROL-LOWERING THERAPY			
(57) Abstract Methods are described for preventing or reducing the risk of a first occurrence of a cardiovascular event using an HMG-CoA reductase inhibitor alone or in combination with another lipid altering agent. Subjects to be treated are those having an average serum total cholesterol level, an average to mildly elevated serum low-density lipoprotein cholesterol level, and a below average serum high-density lipoprotein cholesterol level, with no history of clinically evident coronary disease.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 98/17459

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/365 A61K31/40 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>PROBSTFIELD, JEFFREY L. (1) ET AL: "Results of the primary outcome measure and clinical events from the asymptomatic carotid artery progression study." AMERICAN JOURNAL OF CARDIOLOGY, (1995) VOL. 76, NO. 9, PP. 47C-53C. ISSN: 0002-9149., XP002095840 see abstract</p> <p style="text-align: center;">--- -/--</p>	1-28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

*** Special categories of cited documents :**

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "S" document member of the same patent family

Date of the actual completion of the international search

8 March 1999

Date of mailing of the international search report

01/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

A. Jakobs

INTERNATIONAL SEARCH REPORT

Inter. Appl. Application No
PCT/US 98/17459

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MARCUS A.: "Role of the HMG-CoA reductase inhibitors in the treatment of dyslipidemia An evolutionary review." CARDIOVASCULAR REVIEWS AND REPORTS, (1996) 17/1 (10-11+15-16+21-24+26-27) ISSN: 0197-3118 CODEN: CRRPD4, XP002095841 United States see abstract see page 11, column 2, paragraph 4 - page 22, column 2, paragraph 1 ---	1-28
X	MEISER, B. M. ET AL: "Simvastatin decreases accelerated graft vessel disease (GVD) after heart transplantation in an animal model" TRANSPLANT. PROC. (1993), 25(2), 2077-9 CODEN: TRPPA8;ISSN: 0041-1345, XP002095842 see page 2077, column 1, paragraph 1-4; table 1 see page 2078, column 1, paragraph 1-3 ---	1-28
X	KELLICK K A ET AL: "Outcome monitoring of fluvastatin in a Department of Veterans Affairs lipid clinic." AMERICAN JOURNAL OF CARDIOLOGY, (1995 JUL 13) 76 (2) 62A-64A. JOURNAL CODE: 3DQ. ISSN: 0002-9149., XP002095843 United States see abstract see page 63A, column 1, paragraph 4 - column 2, paragraph 3 ---	1-25, 27, 28
X	MOTOMURA N. ET AL: "HMG-CoA reductase inhibitors in organ transplantation." JOURNAL OF NEPHROLOGY, (1997) 10/2 (68-76). REFS: 127 ISSN: 1121-8428 CODEN: JLNEEL, XP002095844 Italy see abstract; figure 2 see page 69, column 3 - page 70, column 1 see page 72, column 2, paragraph 3 - column 3, paragraph 1 ---	1-28
X	EP 0 528 515 A (MERCK & CO INC) 24 February 1993 see claims 1-4 --- -/--	1-28

INTERNATIONAL SEARCH REPORT

Inter: Jnal Application No
PCT/US 98/17459

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HARIA M. ET AL: "Pravastatin: A reappraisal of its pharmacological properties and clinical effectiveness in the management of coronary heart disease." DRUGS, (1997) 53/2 (299-336). REFS: 267 ISSN: 0012-6667 CODEN: DRUGAY, XP002095845 New Zealand see abstract see page 310 - page 314 see page 323	1-25, 27, 28
X	STEIN E. ET AL: "Cerivastatin, a new potent synthetic HMG Co-A reductase inhibitor: Effect of 0.2 mg daily in subjects with primary hypercholesterolemia." JOURNAL OF CARDIOVASCULAR PHARMACOLOGY AND THERAPEUTICS, (1997) 2/1 (7-16). REFS: 30 ISSN: 1074-2484 CODEN: JCPTFE, XP002095846 United States see abstract see page 7, column 2 - page 8, column 2, paragraph 2; table 8	1-28
X	SOUTHWORTH M R ET AL: "The use of HMG-CoA reductase inhibitors to prevent accelerated graft atherosclerosis in heart transplant patients." ANNALS OF PHARMACOTHERAPY, (1997 APR) 31 (4) 489-91. REF: 32 JOURNAL CODE: BBX. ISSN: 1060-0280., XP002095847 United States see abstract see page 489, column 2, paragraph 6 - page 490, column 2, paragraph 4	1-28
X	PFEIFER M.A. ET AL: "Cholesterol and recurrent events: A secondary prevention trial for normolipidemic patients." AMERICAN JOURNAL OF CARDIOLOGY, (1995) 76/9 (98C-106C). ISSN: 0002-9149 CODEN: AJCDAG, XP000579209 United States see abstract	1-28
X	EP 0 461 548 A (SQUIBB & SONS INC) 18 December 1991 see claims 1-20	1-28
X	EP 0 671 171 A (SQUIBB BRISTOL MYERS CO) 13 September 1995 see claims 1-19	1-28
	--- -/--	

INTERNATIONAL SEARCH REPORT

Int. onal Application No

PCT/US 98/17459

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>US 5 622 985 A (OLUKOTUN ADEOVE Y ET AL)</p> <p>22 April 1997</p> <p>see claims 1-19; examples 1-8</p> <p>-----</p>	1-28

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/17459

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-35
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-35 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: -
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
See FURTHER INFORMATION PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 98 17459

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

In view of the large number of compounds, which are defined by the general definition(s)/formulae used in claims 1-24,27-35, the search had to be restricted for economic reasons. The search was limited to the compounds for which pharmacological data was given and / or the compounds mentioned in the claims, and to the general idea underlying the application. (see Guidelines, chapter III, paragraph 2.3)

Regarding Claim 22: Cerebrovascular events are not commonly known to be part of the group of cardiovascular events

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 98/17459

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0528515	A	24-02-1993	CA 2070149 A JP 6009432 A	05-12-1992 18-01-1994
EP 0461548	A	18-12-1991	AU 7719291 A CA 2042526 A JP 4243839 A US 5622985 A	12-12-1991 12-12-1991 31-08-1992 22-04-1997
EP 0671171	A	13-09-1995	AU 1022795 A CA 2140498 A JP 7206712 A US 5691375 A US 5674893 A	27-07-1995 19-07-1995 08-08-1995 25-11-1997 07-10-1997
US 5622985	A	22-04-1997	AU 5074196 A CA 2172884 A EP 0738512 A JP 8291082 A AU 7719291 A CA 2042526 A EP 0461548 A JP 4243839 A	31-10-1996 20-10-1996 23-10-1996 05-11-1996 12-12-1991 12-12-1991 18-12-1991 31-08-1992